

Influence of **FEED** on average weight (BWT), feed conversion (FC), adjusted feed conversion (AFC), and mortality

Composition		Day 18			Day 50			7-49 Days	
No	Citrus peel lbs/ton	BWT (kg)	FC (kg gain/kg feed)	AFC (kg gain/kg feed)	BWT (kg)	FC (kg gain/kg feed)	AFC (kg gain/kg feed)	Mortality (%)	
1	Control	0.583±0.007	1.252±0.007	1.243±0.007	2.860±0.023	1.987±0.018	1.923±0.014	6.000±1.309	
2	4	0.582±0.006	1.242±0.003	1.231±0.002	2.793±0.025	1.964±0.014	1.915±0.009	4.250±0.796	
3	8	0.581±0.005	1.233±0.007	1.220±0.007	2.824±0.022	1.953±0.016	1.910±0.010	3.250±0.750	
4	16	0.582±0.007	1.244±0.009	1.237±0.009	2.835±0.024	1.959±0.014	1.912±0.011	3.750±1.098	
5	24	0.585±0.006	1.249±0.006	1.241±0.007	2.853±0.022	1.970±0.018	1.919±0.006	4.250±1.278	
6	32	0.577±0.008	1.259±0.009	1.245±0.008	2.890±0.023	2.000±0.025	1.910±0.010	7.250±1.359	
P>F		0.9631	0.1862	0.1262	0.0916	0.4288	0.9335	0.1288	

FIG. 1




FIG. 2

Average Bird Weight And
Feed Conversion




FIG. 3




FIG. 4




FIG. 5




FIG. 6




FIG. 7

Average Body Weights, Gain and
Adjusted Feed Conversion (Days 0 - 15)




FIG. 8

Average Body Weight Day 27 and Adjusted Feed Conversion




FIG. 9